

Tile Sets Generated by Hadamard Submatrices of Fourier Matrices

Troy Wiegand with Faculty Mentor Dr. John Herr
Butler University

What are Hadamard and Fourier Matrices?

A Hadamard Matrix H is a N by N square matrix whose entries all have a complex modulus of one (i.e. 1 unit away from the origin of the complex plane) and $H^*H = HH^* = N \cdot I_N$, where $*$ is the adjoint and I_N is the Identity Matrix of size N by N .

Fourier Matrices are a subclass of Hadamard Matrices that introduce more structure to the matrix. All entries of an M by M Fourier Matrix have the form:

$$f_{jk} = e^{2i\pi \frac{j(k)}{M}},$$

where j is the j th row and k is the k th column (both starting with 0).


Example:

$$\mathcal{F}_4 = \begin{bmatrix} e^{2i\pi \frac{0}{4}} & e^{2i\pi \frac{0}{4}} & e^{2i\pi \frac{0}{4}} & e^{2i\pi \frac{0}{4}} \\ e^{2i\pi \frac{1}{4}} & e^{2i\pi \frac{1}{4}} & e^{2i\pi \frac{2}{4}} & e^{2i\pi \frac{3}{4}} \\ e^{2i\pi \frac{2}{4}} & e^{2i\pi \frac{2}{4}} & e^{2i\pi \frac{4}{4}} & e^{2i\pi \frac{6}{4}} \\ e^{2i\pi \frac{3}{4}} & e^{2i\pi \frac{3}{4}} & e^{2i\pi \frac{6}{4}} & e^{2i\pi \frac{9}{4}} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{bmatrix}$$

Tile Sets over the Integers $\mathbb{Z}$

Tiling by translation is an old problem. Take any finite subset of the integers A . If there exists a sequence T of translations to every element in A without overlap then this sequence T is a Tile Set for A . T can be decomposed into a set of shifts B and a period N_A , so that $T = B \oplus N_A \mathbb{Z}$.

Example:



This A has a B of $\{0, 2\}$. This makes sense because:

- We keep the initial position of the elements of A
 - Shifting by one fails due to overlap.
 - Shifting by two works and creates a consecutive string of 8 numbers.
- Incidentally, the period N_A of $A = \{0, 1, 4, 5\}$ is 8.

The Fuglede Conjecture and the Universal Tiling Conjecture in One Dimension

THEOREM (Fuglede Conjecture (1974)) *A domain Ω admits an operator spectrum if and only if it is possible to tile $\mathbb{R}^d$ by a family of translates of Ω .*

This has been shown to be true in a variety of special cases however it has been shown to be false in both directions in dimensions 3 and higher.

THEOREM (Universal Tiling Conjecture (Dutkay and Jorgensen, 2013)) *Let $p \in \mathbb{N}$. Let $\Gamma := \lambda_0 = 0, \lambda_1, \lambda_{p-1}$ be a subset of $\mathbb{R}$ with p elements. Assume Γ has a spectrum of the form $\frac{1}{p}A$ with $A \subset \mathbb{Z}$. Then for every finite family $A_1, A_2, \dots, A_n$ of subsets of $\mathbb{Z}$ such that $\frac{1}{p}A_i$ is a spectrum for Γ for all i there exists a common tiling subset of $\mathbb{Z}$ such that the set A_i tiles $\mathbb{Z}$ by for all $i \in \{1, \dots, n\}$.*

THEOREM *The following statements are equivalent.*

1. *The Universal Tiling Conjecture is true for all $p \in \mathbb{N}$*
2. *Every bounded Lebesgue measurable spectral set tiles by translations.*

Moreover, if these statements are true and if $\Omega, |\Omega| = 1$, is a bounded Lebesgue measurable set which has a spectrum with period p , then Ω tiles by a subset of $\frac{1}{p}\mathbb{Z}$

Our Problem

Through the findings of Dutkay and Jorgensen we are able to relate these problems to statements about Fourier and Hadamard matrices. We look for two submatrices of the same Fourier Matrix that are formed from the same columns whose row sets do not share a common tiling set. This would yield a counterexample to the Universal Tiling Conjecture, in turn showing that the spectral-tiling direction of the Fuglede Conjecture is false in dimension one.

The Generation of Hadamard Submatrices

We are generating our Hadamard Submatrices through a Mathematica program. It starts with a M by M Fourier matrix and then picks N columns and rows and checks to see if the corresponding submatrix is Hadamard H by examining the entries H^*H . If it ever fails it tries another submatrix. It at worst has an estimated run time of $O((\frac{p}{2})^2 n!)$.

Newman Method

THEOREM *Let $a_1, a_2, \dots, a_k$ be distinct integers with $k = p^\alpha$, p is a prime, α is a positive integer. For each pair $a_i, a_j, i \neq j$, we denote $p^{e_{ij}}$ the highest power of p which divides $a_i - a_j$. The set $a_1, a_2, \dots, a_k$ tessellates or tiles by translation if and only if there are at most α distinct e_{ij} .*

Invoking this theorem is a powerful tool to process a large number of submatrices. We created a program in the language C++ that properly determines if an A can tile given $|A| \leq 15$. This was a powerful tool that helped us look at some of the set of submatrices and develop intuition about tiling.

Coven Meyerowitz (CM) Properties

In 1998 Ethan M. Coven and Aaron Meyerowitz discovered sufficient conditions for a set to tile. Here is some more notation that will be necessary to understand the CM Properties:

- $A(x)$ is a polynomial of the form $\sum_{a \in A} x^a$ and $A(1) = |A|$
- S_A is the set of prime powers s such that s th cyclotomic polynomial $\Phi_s(x)$ divides $A(x)$

Example:

For $A = \{0, 1, 4, 5\}$, $A(x) = x^5 + x^4 + x + 1$ and $S_A = \{2, 8\}$

There are two properties need to be true in order to show that a set tiles:

T1 $A(1)$ needs to equal $\prod_{s \in S_A} \Phi_s(1)$

T2 If $s_1, \dots, s_m \in S_A$ are powers of distinct primes, then $\Phi_{s_1 \dots s_m}$ divides $A(x)$.

THEOREM *If $|A|$ has at most two prime factors, then $A(x)$ satisfies the two CM Properties if and only if A tiles the integers.*

The CM Properties check have been implemented in our C++ program to allow us to check if all the submatrices in our data actually tile.

Determining Period and Tile Set of A

Coven and Meyerowitz also found that the period and tile set of A can be determined through use of $A(x)$ and S_A . The period is defined to be the least common multiple amongst all of the elements of S_A . Finding B is slightly more difficult. First we must define $B(x)$ to be

$$\prod \Phi_{s_B}(x^{t(s_B)}),$$

where s_B range over the prime power factors of $N_A \notin S_A$ and $t(s_B)$ is the largest factor of N_A relative prime to s_B .

Example: Let's look at $A = \{0, 1, 4, 5\}$ again.

Its $S_A = \{2, 8\}$. The period N_A is $lcm(2, 8) = 8$, and $B(x)$ is $\Phi_4(x^1) = x^2 + 1$ because our only s_B is 4 which only has 1 as a relatively prime factor to 8. This confirms that the tiling set $B = \{0, 2\}$ as we showed earlier by hand.

This method is very applicable and useful in testing large numbers of submatrices.

Results

We have found that for all of the Hadamard submatrices from $\mathcal{F}_2$ up to $\mathcal{F}_{18}$ do tile. All of the rows that have a common column set have the same period. However, there can be multiple periods amongst the same size of Hadamard submatrices generated from the same Fourier Matrix.

Further Study

We have developed multiple goals to help guide this project further:

- Determine the tiling set of all our current submatrices.
- Write the Hadamard Submatrix Generator in a less interpreted language that can be run in parallel (C++) to be able to gather more submatrices to analyze
- Increase the capacity of the Tiling Check Program to handle larger submatrices when they are produced.
- Discover a technique to check for set sizes of three or more distinct prime powers.

References

- Bond, Bailey, Glickfield, Alexander, and Herr, John E. (2018). On the Existence of Complex Hadamard Submatrices of the Fourier Matrices.
- Dutkay, Dorin Jorgensen, Palle. (2012). On the Universal Tiling Conjecture in Dimension One. Journal of Fourier Analysis and Applications.
- Dutkay, Dorin Kraus, Isabelle. (2018). On spectral sets of integers. Contemporary Mathematics. 215-234.
- M. Coven, Ethan Meyerowitz, Aaron. (1998). Tiling the Integers with Translates of One Finite Set. Journal of Algebra. 212.
- Newman, David. (1977). Tessellation of integers. Journal of Number Theory - J NUMBER THEOR.